

**Este artículo puede ser usado únicamente para uso personal o académico.  
Cualquier otro uso requiere permiso del auto.**

# Graph and Automata in Arterial Vascular Tree of the Kidney

Aurora Espinoza-Valdez, Ricardo Femat \*

*Laboratorio para Biodinámica y Sistemas Alineales, División de Matemáticas Aplicadas, IPICYT. Apdo. Postal 3-90, Tangamanga c.p. 78231, San Luis Potosí, S. L. P., México*

---

## Abstract

The renal vascular development is not known and occurs through two mechanisms that sometimes overlap: vasculogenesis and angiogenesis. Here we only consider growth through angiogenesis, i.e., the arterial vascular tree of the kidney. There are two types of vascular angiogenesis in development: sprouting and splitting angiogenesis. We study these processes through mathematical tools. The graphs and automatas allow modeling the vascular growth, can generate tree structures by incorporating the physiological laws of the arterial branching. That is, the graph prescribes topology of the vascular tree and the automaton can include the rule of dynamics in the phenomena of vascularization.

**Keywords:** Renal Vasculature, Sprouting and Splitting Angiogenesis, Graph Theory, Automata.

---

## 1 Introduction

The kidney is a highly vascularized organ, the vascular tree of kidney has been the focus of a number of studies (Nordsletten, et. al., 2006; Sequeira and Gomez, 2000; Tomanek, 2001; Zamir, 2001; Zamir and Phipps, 1987). The kidney consists of three trees: arterial, venous, and ureter (Guyton and Hall, 2000). Vasculogenesis and angiogenesis have both been postulated as responsible for the formation of the renal vessels (Tomanek, 2001). The formation of new blood vessels is an important process in embryonic development, vasculogenesis is the formation of blood vessels from endothelial cells precursors.

---

\* Corresponding Author

*Email address:* rfemat@ipicyt.edu.mx (Ricardo Femat ).

This process is predominant during early embryogenesis (Tomanek, 2001). Angiogenesis is defined as the formation of new blood vessels from preexisting vessels. The contribution of each process in the development of the kidney remains to be determined. It is likely, however, that small arterioles and capillaries are formed by vasculogenesis, whereas larger arteries may develop by angiogenesis. Vascular endothelial growth factor (VEGF) plays an important role in renal vascularization. VEGF is probably a crucial parameter in kidney vascularization. The growth of kidney vessels seems to depend on the regulation of VEGF by the tissue oxygen concentration. Both processes depend of VEGF (Tomanek, 2001).

Here, we only consider angiogenesis because we are interested in arterial vascular tree of the kidney. The renal arterial tree is structured as follows: the renal artery had divided into interlobar arteries, arcuate arteries, and interlobular arteries. Angiogenesis is divided in sprouting and splitting. Sprouting angiogenesis is a process in which endothelial cells activate branch out from a existing vessel to produce a new emerging vessel. This process involves degradation of the basement membrane surrounding the endothelial layer, migration of the endothelial cells towards the angiogenic stimulus, proliferation of the endothelial cells, lumen formation and fusion with the preexisting vessel (Patan, 2000). Splitting angiogenesis is a process in which new blood vessels are generated by dividing an existing blood vessel. Splitting may occur as the result of proliferation of endothelial cells inside the vessel lumen and the subsequent fusion and splitting of the vessels (Patan, 2000). In this work, we study the development of arterial vascular tree in the kidney by angiogenesis. The arterial vascular tree of the kidney is modeled using graph theory and automatas. A graph is used to visualize the topology of arterial vascular tree of the kidney providing physiological information into the edges. We can incorporate dynamics of the development in renal arterial tree on the graph and this information can go into edges also. That is, we found the rule of the automata for the dynamics. The paper is organized as follows: Section 2 includes definitions of graph, sprouting and splitting angiogenesis, and tree-automata. Section 3 our results are shown here and the manuscript is closed with some concluding remarks in section 4.

## 2 Definitions

We define a graph to visualize the topology of arterial vascular tree of the kidney with physiological and dynamical information in the edges.

**Definition 1**  $G_R$  is an ordered triple  $(V(G_R), E(G_R), \psi_{G_R})$  that consist of a nonempty set  $V(G_R)$  of vertices, a set  $E(G_R)$  of edges which is disjoint from  $V(G_R)$ , and an incidence function  $\psi_{G_R} : E(G_R) \rightarrow K_{\leq 2}^{V(G_R)}$ , where  $K_{\leq 2}^{V(G_R)}$

is the set of vertices  $\leq 2$ , for each edge is met either of the following two conditions:

- (1)  $\psi_{G_R}$  associates each edge to a subset of  $V(G_R)$  of size two; that is,  $\psi_{G_R}(e) = \{u, v\}$ .
- (2)  $\psi_{G_R}$  associates to each edge, a subset of an element of  $V(G_R)$ ; that is,  $\psi_{G_R}(e) = \{u\}$ .

Let  $G_R$  be a tree, i.e., a connected acyclic graph.  $G_R$  has vertices with oriented edges in such form that of each vertex leave two edges and arrive an edge (the orientation symbolizes blood circulation flow in arteries and ). Thus, given any edges on a bifurcation, these are related as follows

$$i = \begin{cases} \frac{(m-1)}{2} & \text{if subscript } m \text{ is odd} \\ \frac{m-2}{2} & \text{if subscript } m \text{ is even} \end{cases}$$

on which  $i$  denotes the edge generating the other two in the bifurcation and, complementarily, in the tree  $G_R$ , the subscript  $j$  stands for depth of a vessel onto the tree (see Figure 1). Note that each edge includes information about the  $i$ -th pre-existent vessel from which the angiogenesis occurs via sprouting or splitting.




Fig. 1. In  $G_R$  the subscript  $i$  indicate in order to know that edge generates what edge.  $G_R$  has depth  $j$ , each  $j$  is a segment of the tree. The subscript  $t$  ( $t \in \mathbb{N}$ ) indicate the position of the vertex in  $G_R$ .

The tree  $G_R$  has labeled edges, that is, each edge represents a blood vessel and its labeled  $e_{i(j-1)}(s, C_{gf}, l, d, \theta)$  ( $i, j \in \mathbb{N}$ ), i.e., the edge has the physiological information: the parameter  $s$  has the dynamics (depends the process used in the development of the vessel, i.e.,  $a_b$  or  $a_p$ ), concentration of VEGF ( $C_{gf}$ ), length ( $l$ ), diameter ( $d$ ) and angle ( $\theta$ ), see Fig. 2.




Fig. 2. Representation of an arterial bifurcation in  $G_R$  with labeled and oriented edges.

All parameters are important, but  $C_{gf}$  is an essential regulatory factor for both processes of angiogenesis. While  $C_{gf}$  promotes endothelial cell migration, proliferation and tube formation, its effect on splitting angiogenesis has still to be investigated. We define mathematically the processes of sprouting and splitting angiogenesis:

**Definition 2** Let  $a_b$  be a sprouting angiogenesis,  $a_b$  generates a new blood vessel in the edge  $e_{i(j-1)}$ , which is formed by  $k$  ( $k \in \mathbb{N}$ ) endothelial cells.

**Definition 3** Let  $a_p$  be a splitting angiogenesis,  $a_p$  generates two new blood vessels in the edge  $e_{i(j-1)}$ .

We can to define a cellular automata in the tree  $G_R$ , now the set of configurations is  $C = S^{E(G_R)}$  (maps of edges of  $E(G_R)$  to  $S$ ), for every  $e \in E(G_R)$  we denote  $N_e \subseteq E(G_R)$  (finite subset) and is called the set of neighbors of the edge  $e$ , which contains  $e$  and its neighboring. Finally we have for each  $e \in E(G_R)$  a local map  $f_e$  from  $S^{N_e}$  to  $S$  (where  $S^{N_e}$  is the set of maps from  $N_e$  to  $S$ ).

**Definition 4** A  $G_R$ -automata is an ordered pair  $(G_R, A)$  where  $G_R$  is a tree defined by  $(V(G_R), E(G_R), \psi_{G_R})$  as described above and  $A$  is a triple  $(S, N_e, f_e)$ , we have for each element  $e_{i(j-1)} \in E(G_R)$  a finite subset  $N_e$  of  $E(G_R)$  that contains  $e_{i(j-1)}$  (and its neighboring) and is called the set of neighbors of the edge  $e_{i(j-1)}$ . We have a set of values  $S = \{a_b, a_p\}$ . Finally we have for each

### *Sprouting*




Fig. 3. (i) If the new vessel in  $e_{i(j-1)}$  is  $e_{(m-1)j}$ , formed by  $a_b$ , then  $d_{m-1} < d_m$ ,  $\theta_{m-1} > \theta_m$ , and  $d_m = d_i$  or, equivalently, (ii) if the new vessel in  $e_{i(j-1)}$  is  $e_{mj}$ , formed by  $a_b$ , then  $d_m < d_{m-1}$ ,  $\theta_m > \theta_{m-1}$ , and  $d_{m-1} = d_i$  (Gabryś, et. al., 2005; Patan, 2000).

### *Splitting*




Fig. 4. The vessel  $e_{i(j-1)}$  bifurcates in  $e_{(m-1)j}$  and  $e_{mj}$ . Diameters of new vessels are  $d_{m-1} = d_m = \frac{d_i}{2}$  and  $\theta_{m-1} + \theta_m = 75^\circ$  (Gabryś, et. al., 2005; Patan, 2000).

$e_{i(j-1)} \in E(G_R)$  a local map  $f_e : S^{N_e} \rightarrow S$ .

$$f_e(e_{i(j-1)}) = \begin{cases} a_b & si \exists ec \\ a_p & si \nexists ec \end{cases}$$

where  $ec$  is the migration of endothelial cells.  $f_e$  generates each bifurcation in the tree (see Fig. 2).

Here, we only consider  $N_e = e_{i(j-1)}$ , because we do not have experimental data of how it is the dependency with respect to its neighbors of the blood vessels.

We have three possible structures in the bifurcation when the pre-existing edge is  $a_b$ , and three possible structures in the bifurcation when the pre-existing edge is  $a_p$ . All bifurcations have the same probability of  $\frac{1}{3}$ . As a matter of fact,

$$\begin{aligned} \text{If } e_{i(j-1)} \text{ is } a_b, \Rightarrow \text{ can be generated the bifurcations} &= \begin{cases} a_b a_b, \\ a_b a_b, \text{ or} \\ a_p a_p \end{cases} \\ \text{or} & \\ \text{If } e_{i(j-1)} \text{ is } a_p, \Rightarrow \text{ can be generated the bifurcations} &= \begin{cases} a_b a_p, \\ a_p a_b, \text{ or} \\ a_p a_p \end{cases} \end{aligned}$$

The local maps  $f_e$  induce a global map  $F$  from the set of configurations to itself as follows. Let  $c$  be a configuration, we define the configuration  $F(c)$  by:

$$\forall e_{i(j-1)} \in E(G_R) \quad F(c)(e_{i(j-1)}) := f_e(c|_{N_e}),$$

where  $c|_{N_e}$  is the restriction of  $c$  to the set  $N_e$ .

### 3 Results

Our results are discussed in context of the development of vascular tree in the kidney from the angiogenesis. By definition of sprouting and splitting angiogenesis, we have that these processes are different and the most important different is that  $a_b$  has migration of endothelial cells whereas  $a_p$  does not have.

**Axiom 1**  $a_b \neq a_p$ .

According to experimental studies in the vascular tree of the kidney, we conclude that the renal arterial tree reaches the depth of the interlobular arteries (Guyton and Hall, 2000).

**Proposition 1** *Let  $G_R$ -automata be an arterial vascular tree of the kidney formed by  $a_b$  and  $a_p$ , depth of the tree  $G_R$  is  $0 \leq j \leq 9$ .*

**Proof** Let  $G_R$ -automata be an arterial vascular tree of the kidney, which is formed by  $a_b$  and  $a_p$ . The arterial vascular tree of the kidney is structured: the renal artery is branched in interlobar arteries, arcuate arteries, interlobular arteries which are formed by bifurcation (Guyton and Hall, 2000). In fact, a vascular tree of the kidney until the interlobular arteries can structured as follows: renal artery, segment 0; interlobar arteries, segments 1-2; arcuate arteries, segments 3-4; interlobular arteries, segments 5-9. Hence, we have that the depth of  $G_R$  is  $0 \leq j \leq 9$ .

We have that  $a_b$  generates only one new vessel while  $a_p$  generates two vessels, we can associate on each branching point one vertex  $v_t$ , only two vessel are found after each vertex (see Fig. 1).

**Theorem 1** *If the arterial vascular tree of the kidney is developed through  $a_b$  and  $a_p$ , each segment has an even number of blood vessels ( $b_v$ ).*

**Proof** The arterial vascular tree of the kidney is developed obeying the following steps indistinctly if  $a_b$  or  $a_p$  occur on each vertex: For the segment  $j = 0$ , the renal artery is the unique branch on the vasculature. As vascular development occurs, for the segment  $j = 1 \exists 2 b_v$ . For the segment  $j = 2 \exists 4 b_v$ , that is, we have  $2 \cdot 2 = 2^2$ . Then, inductively, we have that for  $j = n$ ,  $n \in \mathbb{N}$ ,  $\exists 2^n b_v$ . Hence, for the  $n$ -th segment, we have  $2 \cdot 2^{n-1} = 2^n b_v$ .

Therefore, if the arterial vascular tree of the kidney  $G_R$ -automata is developed by means of  $a_b$  and  $a_p$ , each segment  $0 < j \leq 9$  (by proposition 1) has an even number of  $b_v$ .

**Corollary 1** *All vertices in the arterial vascular tree of the kidney  $G_R$  has degree 3, i.e.,  $\forall v \in V(G_R) \text{ deg}_{G_R}(v) = 3$ .*

Initially for  $j = 0$  in the  $G_R$ -automata, there exists the initial configuration  $c_0 = \{e_{00}\}$ , which is the initial condition (the renal artery). In general, for the segment  $j = n \exists$  the configuration  $c_n$ , which have exactly  $2^n$  labeled edges by  $f_e$ . Then, we have

$$c_{n+1} = \{f_e(c_n(e_{1n}), c_n(e_{2n}), \dots, c_n(e_{2^n n}))\}$$

where  $f_e(c_n(e_{2^n n}))$  generates each bifurcation.

All configuration  $c_j$  has  $2^j$  edges with  $2^j = l + k$  vessels, where  $l$  is the number of labeled edges by  $a_b$  and  $k$  is the number of labeled edges by  $a_p$ :

- If  $l$  and  $k \neq 0$ ,  $\exists (2l)(3k)$  possible configurations for generate the next configuration  $c_{j+1}$ .
- If  $l = 0$  or  $k = 0$ ,  $\exists 3k$  or  $2l$  possible configurations for generate the next configuration  $c_{j+1}$ .  $l = 0$  and  $k \neq 0$  is greater in number of possible configurations ( $c_{n+1}$ ) that when  $l \neq 0$  and  $k = 0$ .
- When  $l = k$ , we have the number maximum of possible configurations for the next configuration  $c_{j+1}$ .

For each  $c_j \exists (2l)(3k)$  possible configurations for generate the next configuration  $c_{j+1}$ .

**Proposition 2** *If in the configuration  $c_j \exists$  labeled edges with  $a_b$  and  $a_p$ , it is possible generate the configuration  $c_{j+1}$  with all labeled edges by  $a_p$ .*



**Proof**  $\exists$  labeled edges with  $a_b$  and  $a_p$  in  $c_n$ . Then, we have that the bifurcation of  $a_b$  (pre-existing vessel) can have labeled edges by  $a_p$  and the bifurcation of  $a_p$  (pre-existing vessel) can have labeled edges by  $a_p$  (see Figure XX). Consequently it is possible generate the configuration  $c_{j+1}$  will all labeled edges  $a_p$ .

When we have all labeled edges by  $a_b$  or  $a_p$ , then it is impossible to generate the next configuration  $c_{j+1}$  with all labeled edges by  $a_b$ , due to the presence of  $a_p$  in  $c_j$ .

**Proposition 3** *If in the configuration  $c_n$  all edges are  $a_b$ , it is possible generate the next configuration  $c_{j+1}$  with all labeled edges by  $a_b$  or  $a_p$ .*

**Proof** In the configuration  $c_n$  all edges are  $a_b$ , we have that the bifurcation of  $a_b$  (pre-existing vessel) can have labeled edges by  $a_b$  or  $a_p$ , and the bifurcation of  $a_p$  (pre-existing vessel) can have labeled edges by  $a_b$  or  $a_p$  too. Then, it is possible generate the configuration  $c_{n+1}$  will all labeled edges by  $a_b$  or  $a_p$ .

The degree of diametral asymmetry on a bifurcation is expressed by the index:  $\alpha = \frac{d_{m-1}}{d_m}$ , where  $0 < \alpha \leq 1$  and diameters  $d_{m-1}$  and  $d_m$  are related to discussion on the Figure XX. The index is a measure of the asymmetry of the two diameters at an arterial bifurcation. As  $\alpha = 1$ , both diameters are equal (i.e.,  $d_{m-1} = d_m$ ), and, oppositely, one of the two diameters is larger than the other (i.e.,  $d_{m-1} < d_m$ ) as  $\alpha < 1$ .

**Theorem 2** *If there exist  $a_b$  and  $a_p$  in the developed of arterial vascular tree of the kidney, tree is asymmetric.*

**Proof**  $\exists a_b$  and  $a_p$  in  $G_R$ , we have that if a new vessel is formed in the bifurcation by  $a_b$ , which implies that  $d_{m-1} < d_m$ , the index of diametral asymmetry is  $\alpha < 1$ . If the two new vessels are formed in the bifurcation by  $a_p$ , which implies  $d_{m-1} = d_m$ , the index of diametral asymmetry is  $\alpha = 1$ . Hence,  $\alpha$  is not constant in developing the arterial vascular tree of the kidney.

We observe that the sprouting angiogenesis contributes to an increase of complexity in arterial vascular tree of the kidney.

## 4 Conclusions

We studied the development of arterial vascular tree of the kidney using the graph theory and automatas. We define a  $G_R$ -automata to represent the development of arterial vascular tree of the kidney, which is generated through angiogenesis. Since there exist two types of angiogenesis: sprouting and splitting, we analyzed the renal arterial vasculature development through of these

processes. The development dynamics in the arterial vascular tree of the kidney on  $G_R$  and the physiological information can go into edges. That is, we found the rule of the automata for the dynamics.

There exists six possible bifurcations from sprouting and splitting in the arterial vascular tree. By definition sprouting and splitting are two different processes in the development of the  $G_R$ -automata. Then, we have that  $a_b$  generates one whereas  $a_p$  generates two blood vessels, we conclude that the angiogenesis generates only bifurcations in the  $G_R$ -automata. Consequently, each segment  $j > 0$  has an even number of blood vessels ( $j = 0$  has the renal artery), i.e.,  $\forall v \in V(G_R) \text{ deg}_{G_R}(v) = 3$ . The arterial vascular tree of kidney has a depth until the interlobular arteries, and we determine the depth of the  $G_R$ -automata is  $0 < j \leq 9$ . For each configuration  $c_j$  there exists  $2^j = (2l)(3k)$  (where  $l$  is the number of labeled edges by  $a_b$  and  $k$  is the number of labeled edges by  $a_p$ ) possible configurations for generate the next configuration  $c_{j+1}$ . If all edges are  $a_b$  in  $c_j$ , it is possible generate the configuration  $c_{j+1}$  with all edges labeled by  $a_b$  or  $a_p$ , whereas if all edges are  $a_b$  and  $a_p$  in  $c_j$ , it is possible generate the configuration  $c_{j+1}$  with all labeled edges by  $a_p$ . Additionally, we conclude that the tree  $G_R$  is asymmetric when the arterial vascular tree of the kidney is developed by  $a_b$  and  $a_p$ , which is consistent with experimental data.

## References

- Gabryś E., Rybaczuk M., and Kedzia A. (2005). Fractal models of circulatory system. Symmetrical and asymmetrical approach comparison. *Chaos, Solutions and Fractals* **24**:707-715.
- Guyton A. C. and Hall J. E. (2000). Tratado de Fisiología Medica. Décima edición. Eritorial McGraw-Hill (In Spanish).
- Nordsletten D. A., Blackett S., Bentley M. D., Ritman E. L., and Smith N. P. (2006). Structural morphology of renal vasculature. *American Journal of Physiology-Heart and Circulatory Physiology*. **291**:H296-H309.
- Patan S. (2000). Vasculogenesis and angiogenesis as mechanisms of vascular network formation, growth and remodeling. *J. Neuro-Oncology*. **50**:1-15.
- Sequeira López M. L. S. y Gomez R. A. (2000). Desarrollo de la vasculatura renal. *Medicina*. **60**:694-698 (In Spanish).
- Tomake R. J. editor (2001). Assembly of the vasculature and its regulation. Berlin:Birkhauser.
- Zamir, M. (2001). Arterial branching within the confines of fractal L-system formalism. *Jornual of General Physiology*. **118**:267-275.
- Zamir M. and Phipps S. (1987). Morphometric analysis of the distributing vessels of the kidney. *Canadian Journal of Physiology and Pharmacology* **65**:2433-2440.