

This is an Accepted Manuscript of an article published by Taylor & Francis in Journal of Environmental Science and Health, Part B: Pesticides, Food Contaminants, and Agricultural Wastes on 14 Oct 2019, available online: <https://doi.org/10.1080/03601234.2019.1674103>

Cite this article as:

Victor E. Balderas-Hernández, José T. Ornelas-Salas, Ana P. Barba de la Rosa & Antonio De Leon-Rodriguez (2020) Diminution of migration of phthalic acid esters in tequila beverage by the year of production, Journal of Environmental Science and Health, Part B, 55:2, 148-154, DOI: [10.1080/03601234.2019.1674103](https://doi.org/10.1080/03601234.2019.1674103)

© 2020 This manuscript version is deposited under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.

Diminution of Migration of Phthalic Acid Esters in Tequila Beverage by the Year of Production

Victor E. Balderas-Hernández¹, José Tomás Ornelas-Salas², Ana P. Barba de la Rosa¹ and Antonio De Leon-Rodriguez ^{1*}

¹Molecular Biology Department, Institute for Scientific and Technological Research of San Luis Potosí, Camino a la Presa San José 2055, CP 78216, San Luis Potosí, México.

²Faculty of Chemical Sciences, Autonomous University of Guadalajara, Av. Patria 1201, Lomas del Valle 3^a Sección, Zapopan, Jalisco, C.P. 45129, México.

*Corresponding author; Phone: +52 (444) 834-2000; Fax: +52 (444) 8342010. E-mail:

aleonr@ipicyt.edu.mx, aleonr@me.com

ABSTRACT

The presence of diethyl-phthalate (DEP), dibutyl-phthalate (DBP), butylbenzyl-phthalate (BBP), diethylhexyl-phthalate (DEHP) and diisononyl-phthalate (DINP) was determined in 295 tequila samples. They were grouped by age of maturation and year of production. Gas Chromatography coupled with Mass Spectrometry was used for identification and quantification. The results showed 65 (22%) samples were phthalate free. DEP (0.13-0.27 mg/kg), BBP (0.05-2.91 mg/kg) and DINP (1.64-3.43 mg/kg) were detected in 11 (3.73%), 37 (12.54%) and 5 (1.69%) samples, respectively, but these concentrations did not exceed the limits admitted for alcoholic beverages. DBP (0.01- 2.20 mg/kg) and DEHP (0.03-4.64 mg/kg) were detected in 96 (32.54%) and 224 (75.93%) samples, from them only 10 (3.39%) and 15 (5.08%) samples exceed the limits admitted for alcoholic beverages and they were few tequilas produced in the year 2014 or before. All tequilas produced in 2015 and after, satisfied the international standards for these compounds. We concluded that a moderate consumption of tequila does not pose a health risk regarding the tolerable daily intake of phthalates according to the international standards for no fat meals and distilled alcoholic beverages.

Keywords: Alcoholic beverages, endocrine disruptor, phthalate, tequila, spirits.

1. INTRODUCTION

Phthalic acid esters or phthalates (Fig. 1) are plasticizer substances widely used in several industrial processes and they have been detected in toys, food packaging, medical materials and other articles ^[1–3]. Phthalates are not chemically bonded to the plastic and they can migrate and contaminate food and beverages during manufacturing or storage ^[4–7]. Special attention has been paid to the presence of phthalates in food and beverages as they have been referred to as substances with endocrine disruptor and cytotoxicity activity ^[1,8]. Experimental phthalate exposition using rats has been shown to cause anti-androgenic effects such as cryptorchidism, diminution of testosterone levels and infertility ^[9–11]. Studies on human exposition to phthalates and their metabolites have been correlated with DNA damage in adult sperm as well as genital malformation among male infants with prenatal exposition ^[12,13]. Phthalates have been related to cancer development; for instance, dibutyl phthalate and benzyl butyl phthalate increase the *in vitro* and *in vivo* proliferation of human breast cancer cell-lines ^[14]. International safety authorities have regulated the concentration limits of certain phthalates in consumer products ^[1]. In the case of foods and beverages, international regulations ^[15,16] have established the permissible limits for some phthalates as shown in Table 1.

Tequila is the main alcoholic beverage produced in Mexico from the distillation and fermentation of sugars from *Agave tequilana* Weber var. *azul*. Tequila has the denomination of origin and exclusive trademarks with worldwide recognition. Tequila is produced in factories with steel fermenters and distilling devices, and it is classified as white or silver tequila when it is distilled and bottled after the alcoholic grade is fixed to 38°alc. vol.; aged, extra aged or ultra aged, for tequila matured in oak casks for at least 3, 12, and 36 months, respectively in agreement with the Mexican Official norm NOM-006-SCFI-2012 ^[17,18].

The aim of this work was to determine the phthalate content in a tequila collection of 295 samples produced from 2013-2018 by gas chromatography coupled to mass spectrometry. This analytical technique has been adopted for the routine analysis of phthalates in alcoholic beverages and biological samples such as serum and urine ^[19–21].

2. MATERIALS AND METHODS

2.1. Chemicals

Diethyl-phthalate (DEP), dibutyl-phthalate (DBP), butylbenzyl-phthalate (BBP), diethylhexyl-phthalate (DEHP), diisononyl- phthalate (DINP) and other chemicals were purchased from Sigma-Aldrich with purity up of 99.9%. Methanol (HPLC grade) was used to wash all glass material to eliminate lab phthalate contamination, which was also confirmed by analysis as described below.

2.2. Samples Collections

295 tequila samples (glass bottled) produced from 2013 to 2018 were analyzed, from them 185 were white, 48 aged, 36 extra aged and 26 ultra aged tequilas (Table 2). Each sample was processed as described previously by Gonzalez-Castro et al.^[2] and filtrated using 0.22 µm filters (Millipore).

2.3. Instrumental Analysis

Samples were analyzed by a Gas Chromatograph coupled to a Mass Spectrometry detector 7820A/5977E System (Agilent Technologies) with an automatic injector 7683 Series (Agilent Technologies) using a capillary column HP5-MS (Agilent Technologies) 30 m long, 250 µm inner diameter and 0.25 µm-film thickness. The injector temperature was 250°C. The temperature program

started at 60°C for 1 min, and it was increased at a rate of 20°C/min until 220°C. The temperature was held at 220°C for 1 min, then increased at a rate of 5°C/min to 280°C and held at 280°C for 4 min. The ionization potential was 70 eV and a scan function with 45-300 m/z for identification and SIM mode for quantification ^[2]. Compounds were identified by comparing their mass spectra with those obtained in the NIST 14 library (Gaithersburg, MD, USA) from the MS database, and by using commercial standards (Table 3).

2.4. Method Validation

Detection and quantification limits were measured by the signal/noise method and by using the standard deviation of the response and the slope ^[22]. The linear response of detectors was determined by straight-line calibration curves from the analysis of five standards with increasing the phthalate concentrations (Table 4) ^[23].

3. RESULTS AND DISCUSSION

A summary of phthalates detected in the tequila samples is shown in Table 5 and Fig. 2. As noted 65 samples (22%) were phthalate free, 112 samples (38%) contained one phthalate (DEHP), 99 samples (33.6%) presented two phthalates (DEHP and DBP), 13 tequila samples (4.4%) contained three phthalates (DEHP, DBP and BBP), and six samples (2%) presented four phthalates (DBP, BBP, DEHP and DEP or DINP). None tequila samples showed five phthalate simultaneously.

Tequila were grouped with respect to the age of maturation as White (n=185), Aged (n=48), Extra aged (n=36), and Ultra aged (n=26) (Table 2). From them, DEP (0.13-0.27 mg/kg), BBP (0.05-2.91 mg/kg) and DINP (1.64-3.43 mg/kg) were detected in 11, 37 and 5 samples, respectively. In all cases, phthalate

concentration did not exceed the admitted limit for alcoholic beverages. DBP was detected in 96 tequilas from the total samples representing the 32.5%, but only 3.39% exceeded the limit of 0.3 mg/kg. Meanwhile 224 samples (75.9%) contained DEHP and 5.08 % exceeded the permissible limit of 1.5 mg/kg (Table 6).

Since DEHP was the most frequent phthalate detected, special attention was paid to this phthalate. Fig. 3 shows DEHP content in tequila samples grouped by age of maturation and year of production. As noted White tequilas presented DEHP below 0.47 ± 0.35 mg/kg or less. Only aged tequila produced in 2013 exceeded the permitted limit with an average concentration of 2.2 ± 1.86 mg/kg, whereas the subsequent years the DEHP concentration dropped to 0.08 ± 0.02 mg/kg in 2018 samples. Extra aged tequilas from 2013 and 2014 showed DEHP of 0.97 ± 0.81 and 1.1 ± 1.4 mg/kg, respectively, indicating that some samples were out of the permitted limit, but in the subsequent years, the DEHP concentration was reduced until 0.17 ± 0.04 mg/kg. Ultra aged tequilas produced from years 2014 to 2018 showed DEHP from 1.2 ± 0.41 to 0.19 ± 0.04 mg/kg. The important reduction of DEHP content in tequila is evidenced in the drop from 1.49 ± 1.58 to 0.14 ± 0.06 mg/kg from 2013 to 2018 total production, as a result of the improvement on the factories, quality control enhancement and reduction in the use of synthetic materials during the beverage elaboration.

DEHP is still the most used plasticizer and also the most studied phthalate; it has been detected in quotidian materials including food and alcoholic and non-alcoholic beverages. Some authors refer to it as an “omnipresent contaminant” [24]. Ye et al. reported that DBP and DEHP were the main phthalates found in three bottles of Chinese beer, and the total concentrations were between 6.22 and 7.76 $\mu\text{g/L}$ [25]. Migration tests revealed that the high content of DEHP incorporated in polyvinyl chloride gaskets in the lids could be a potential source of phthalate contamination in bottled beers during transportation

and storage. Studies on DEHP metabolism in humans showed that the major metabolite in serum was mono 2-ethylhexyl phthalate (MEHP), whereas the major metabolite in urine was 5OH-MEHP, 5oxo-MEHP and MEHP ^[24]. These metabolites are shown after short-term oral-exposition to DEHP, whereas mono 2-ethyl-5-carboxypentyl phthalate (5cx-MEPP) and mono [2-(carboxymethyl)hexyl] phthalate (2cx-MMHP) are also found after long-term oral exposure ^[26].

Other phthalates have been detected in alcoholic beverages, for instance Russo et al. ^[27] analyzed 6 commercial samples of Italian wine; they found 7.3 to 23 µg/L DBP, 0.1 to 7 µg/L BBP, and 3.1 to 16 µg/L DEHP. Cinelli et al. ^[21] found 0.0333 to 0.3124 mg/L of DBP, 0.046 to 0.135 mg/L BBP and 0.0115 to 0.0266 mg/L DEHP in 11 samples of Italian wine. Duca et al. ^[28] reported that 8% of bottled wine from a total of 2000 samples produced in the Republic of Moldova had DBP >0.3 mg/kg, which exceeds the limit established by the international regulation. Montevecchi et al. ^[29] studied the repartition of phthalates during distilling of wine for spirit production such as brandy; they reported values up 0.62 mg±0.05 and 0.47 ±0.04 for DBP and DEHP, respectively. They concluded it is very difficult to suggest modification of the distillation process without impairing the quality of brandies. Montevecchi et al. ^[30] determined phthalates by GC/MS in brandy samples from 1987 to 2014. They reported values up to 0.4 mg/kg DBP and 4.18 mg/kg DEHP, for samples produced in 1987 and 1989, respectively, whereas brandies produced in 2014, values were reduced to 0.03 mg/kg DBP and 0.3 mg/kg DEHP, respectively. Jurica et al. analyzed seven phthalates in samples during the plum spirit production and in the final product (plum is a spirit manufactured by registered producers from five European countries), they reported values of 0.822 mg/kg DBP and 1.638 mg/kg DEHP, which exceeded the limits established by the international regulations, but they concluded that a moderate daily consumption of plum spirit does not pose a health risk regarding the Tolerable Daily Intake of BBP, DEHP, and DBP ^[31].

4. CONCLUSIONS

The great majority of samples analyzed here did not exceed the limit accepted by the international regulations, but some samples of tequila produced in 2014 or previous years exceeded the allowable limits for DBP or DEHP, but beverages produced from 2015 to 2018 do satisfied fully the permissible limits of the international standards authorized for beverages. Our results suggest the risk of human exposure to phthalate by consuming tequila is really low and insignificant. Whit this, we concluded that tequila is a safety alcoholic beverage, however, routine monitoring in each lot of beverages produced and commercialized must be mandatory to guarantee that the beverages are safe for human consumption. On the other hand, monitoring and analysis at every stage of the production process including raw materials, fermentation and packing is strongly recommended in order to detect and discard possible phthalate contamination points.

5. ACKNOWLEDGMENTS

We thank partial funding of CONACyT Problemas Nacionales 4601, Rodrigo Juarez by technical support and Graham Tippet for the English language revision.

6. REFERENCES

- [1] Kamrin, M.A. Phthalate Risks, Phthalate Regulation, and Public Health: A Review. *J. Toxicol. Environ. Heal. B-Critical Rev.* **2009**, *12*(2), 157–174.
- [2] Gonzalez-Castro, M.I.; Olea-Serrano, M.F.; Rivas-Velasco, A.M.; Medina-Rivero, E.; Ordonez-

- Acevedo, L.G.; Leon-Rodriguez, A. De. Phthalates and Bisphenols Migration in Mexican Food Cans and Plastic Food Containers. *Bull. Environ. Contam. Toxicol.* **2011**, *86*(6), 627–631.
- [3] Gonzalez-Castro, M.I.; Ordonez-Acevedo, L.G.; Herrera-Vazquez, Y.; Ramirez-Telles, J.A.; Leon-Rodriguez, A. De. Analysis of Phthalic Acid, Bisphenol a and Bisphenol a Dimethacrylate in Mexican Food Cans by Hplc with Evaporative Light Scattering Detector. *Acta Aliment.* **2013**, *42*(2), 229–235.
- [4] Bosnir, J.; Puntaric, D.; Galic, A.; Skes, I.; Dijanic, T.; Klaric, M.; Grgic, M.; Curkovic, M.; Smit, Z. Migration of phthalates from plastic containers into soft drinks and mineral water. *Food Technol. Biotechnol.* **2007**, *45*(1), 91–95.
- [5] Li, Z.K.; Xue, F.; Xu, L.G.; Peng, C.F.; Kuang, H.; Ding, T.; Xu, C.L.; Sheng, C.Y.; Gong, Y.X.; Wang, L.B. Simultaneous Determination of Nine Types of Phthalate Residues in Commercial Milk Products Using HPLC-ESI-MS-MS. *J. Chromatogr. Sci.* **2011**, *49*(4), 338–343.
- [6] Fasano, E.; Bono-Blay, F.; Cirillo, T.; Montuori, P.; Lacorte, S. Migration of phthalates, alkylphenols, bisphenol A and di(2-ethylhexyl)adipate from food packaging. *Food Control* **2012**, *27*(1), 132–138.
- [7] Jeddi, M.Z.; Rastkari, N.; Ahmadkhaniha, R.; Yunesian, M. Concentrations of phthalates in bottled water under common storage conditions: Do they pose a health risk to children? *Food Res. Int.* **2015**, *69*, 256–265.
- [8] Halden, R.U. Plastics and Health Risks. *Annu. Rev. Public Heal.* Vol 31 **2010**, *31*, 179–194.
- [9] Fisher, J.S.; Macpherson, S.; Marchetti, N.; Sharpe, R.M. Human “testicular dysgenesis syndrome”: a possible model using in-utero exposure of the rat to dibutyl phthalate. *Hum. Reprod.* **2003**, *18*(7), 1383–1394.
- [10] Jarfelt, K.; Dalgaard, M.; Hass, U.; Borch, J.; Jacobsen, H.; Ladefoged, O. Antiandrogenic effects in male rats perinatally exposed to a mixture of di(2-ethylhexyl) phthalate and di(2-ethylhexyl) adipate. *Reprod. Toxicol.* **2005**, *19*(4), 505–515.

- [11] Foster, P.M.D. Disruption of reproductive development in male rat offspring following in utero exposure to phthalate esters. *Int. J. Androl.* **2006**, 29(1), 140–147.
- [12] Swan, S.H.; Main, K.M.; Liu, F.; Stewart, S.L.; Kruse, R.L.; Calafat, A.M.; Mao, C.S.; Redmon, J.B.; Ternand, C.L.; Sullivan, S.; Teague, J.L.; Team, S.F.F.R. Decrease in anogenital distance among male infants with prenatal phthalate exposure. *Environ. Health Perspect.* **2005**, 113(8), 1056–1061.
- [13] Hauser, R.; Meeker, J.D.; Singh, N.P.; Silva, M.J.; Ryan, L.; Duty, S.; Calafat, A.M. DNA damage in human sperm is related to urinary levels of phthalate monoester and oxidative metabolites. *Hum. Reprod.* **2007**, 22(3), 688–695.
- [14] Hsieh, T.H.; Tsai, C.F.; Hsu, C.Y.; Kuo, P.L.; Hsi, E.; Suen, J.L.; Hung, C.H.; Lee, J.N.; Chai, C.Y.; Wang, S.C.; Tsai, E.M. N-Butyl Benzyl Phthalate Promotes Breast Cancer Progression by Inducing Expression of Lymphoid Enhancer Factor 1. *PLoS One* **2012**, 7(8).
- [15] AVIS 03-2014. Avis approuvé lors de la séance plénière du Comité scientifique du 21/02/2014. Comité Scientifique De L’agence Fédérale Pour La Sécurité De La Chaine Alimentaire. Available at http://www.afsca.be/comitescientifique/avis/_documents/AVIS03-2014_FR_DOSSIER20. .
- [16] European Community. Commission Regulation (EU) No 10/2011 of 14 January 2011 on plastic materials and articles intended to come into contact with food (Text with EEA relevance) 112. 1-89. Available at <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2011:012:0001:00>. .
- [17] Leon-Rodriguez, A. De; Gonzalez-Hernandez, L.; la Rosa, A.P.B. De; Escalante-Minakata, P.; Lopez, M.G. Characterization of volatile compounds of mezcal, an ethnic alcoholic beverage obtained from Agave salmiana. *J. Agric. Food Chem.* **2006**, 54(4), 1337–1341.
- [18] Leon-Rodriguez, A. De; Escalante-Minakata, P.; Jimenez-Garcia, M.I.; Ordonez-Acevedo, L.G.; Flores, J.L.F.; la Rosa, A.P.B. de. Characterization of Volatile Compounds from Ethnic Agave Alcoholic Beverages by Gas Chromatography-Mass Spectrometry. *Food Technol. Biotechnol.* **2008**, 46(4), 448–455.

- [19] Gao, J.; Yang, C.; Ye, C.; Li, X. [Determination of trace phthalates in beer by gas chromatography coupled with solid-phase microextraction using a calix[6] arene fiber]. *Se Pu* **2009**, 27(3), 356–358.
- [20] Kondo, F.; Ikai, Y.; Hayashi, R.; Okumura, M.; Takatori, S.; Nakazawa, H.; Izumi, S.; Makino, T. Determination of Five Phthalate Monoesters in Human Urine Using Gas Chromatography-Mass Spectrometry. *Bull. Environ. Contam. Toxicol.* **2010**, 85(1), 92–96.
- [21] Cinelli, G.; Avino, P.; Notardonato, I.; Centola, A.; Russo, M. V. Rapid analysis of six phthalate esters in wine by ultrasound-vortex-assisted dispersive liquid-liquid micro-extraction coupled with gas chromatography-flame ionization detector or gas chromatography-ion trap mass spectrometry. *Anal. Chim. Acta* **2013**, 769, 72–78.
- [22] Long, G.L.; Winefordner, J.D. Limit of Detection. *Anal. Chem.* **1983**, 55(7), A712-.
- [23] Thompson, M.; Ellison, S.L.R.; Wood, R. Harmonized guidelines for single-laboratory validation of methods of analysis - (IUPAC technical report). *Pure Appl. Chem.* **2002**, 74(5), 835–855.
- [24] Koch, H.M.; Bolt, H.M.; Angerer, J. Di(2-ethylhexyl)phthalate (DEHP) metabolites in human urine and serum after a single oral dose of deuterium-labelled DEHP. *Arch. Toxicol.* **2004**, 78(3), 123–130.
- [25] Ye, C.W.; Gao, J.; Yang, C.; Liu, X.J.; Li, X.J.; Pan, S.Y. Development and application of an SPME/GC method for the determination of trace phthalates in beer using a calix[6]arene fiber. *Anal. Chim. Acta* **2009**, 641(1–2), 64–74.
- [26] Koch, H.M.; Bolt, H.M.; Preuss, R.; Angerer, J. New metabolites of di(2-ethylhexyl)phthalate (DEHP) in human urine and serum after single oral doses of deuterium-labelled DEHP. *Arch. Toxicol.* **2005**, 79(7), 367–376.
- [27] Russo, M. V; Notardonato, I.; Cinelli, G.; Avino, P. Evaluation of an analytical method for determining phthalate esters in wine samples by solid-phase extraction and gas chromatography coupled with ion-trap mass spectrometer detector. *Anal. Bioanal. Chem.* **2012**, 402(3), 1373–1381.

- [28] Duca, G.; Sturza, R.; Gaina, B.; Lazacovici, D. Analysis of the phthalate content levels in wine products. *Chem. J. Mold.* **2011**, *6*(2), 65–69.
- [29] Montevecchi, G.; Masino, F.; Pascale, N. Di; Vasile Simone, G.; Antonelli, A. Study of the repartition of phthalate esters during distillation of wine for spirit production. *Food Chem* **2017**, *237*, 46–52.
- [30] Montevecchi, G.; Masino, F.; Zanasi, L.; Antonelli, A. Determination of phthalate esters in distillates by ultrasound-vortex-assisted dispersive liquid-liquid micro-extraction (USVADLLME) coupled with gas chromatography/mass spectrometry. *Food Chem* **2017**, *221*, 1354–1360.
- [31] Jurica, K.; Brčić Karačonji, I.; Lasić, D.; Vukić Lušić, D.; Anić Jurica, S.; Lušić, D. Determination of phthalates in plum spirit and their occurrence during plum spirit production. *Acta Aliment.* **2016**, *45*(1), 141–148.

7. FIGURE CAPTIONS

Fig. 1. Structure of common phthalates. The structures of the phthalates: a) Diethyl phthalate (DEP), b) Dibutyl phthalate (DBP) structure, c) Benzyl butyl phthalate (BBP) structure, d) Bis (2-ethylhexyl) phthalate (DEHP) structure, and e) diisononyl phthalate.

Fig. 2. Percentage of beverages containing no phthalate, one or more phthalate simultaneously. 0: Phthalate free; 1: DEHP; 2: DEHP and DBP; 3: DEHP, DBP and BBP; 4: DEHP, DNP, BBP and DEP or DINP; 5: DEHP, DNP, BBP, DEP and DINP.

Fig. 3. DEPH content in tequila samples grouped by age of maturation and year of production.

8. FIGURES (DIAGRAMS) AND TABLES

Table 1 Phthalates concentrations permitted in foods according to international standards for no fat meals including distilled alcoholic beverages.

Abbreviation	Name	Limit premised (mg/kg)
DEP	Diethyl phthalate	Not reported
DBP	Dibutyl phthalate	0.3
BBP	Benzyl butyl phthalate	30.0
DEHP	Di-2-ethyl-hexyl phthalate	1.5
DINP	Diisononyl phthalate	9.0

Table 2. Summary of tequila samples (n) analyzed in this work.

Year	White	Aged	Extra aged	Ultra aged	Total
2013	4	8	3	0	15
2014	86	22	10	6	124
2015	49	7	5	3	64
2016	18	5	6	7	36
2017	24	3	8	7	42
2018	4	3	4	3	14
Total	185	48	36	26	295

Table 3 Summary of chemical characteristics of the phthalate analyzed.

Chemical	CAS number	Retention time (min)	Ions for identification	Ion for quantification	Molecular weight (g/mol)
DEP	84-66-2	8.72	149, 177, 76, 105	149	222,24
DBP	84-74-2	11.29	149, 104, 76, 223	149	278,34
BBP	85-68-7	15.41	149, 91, 206, 65	149	312.36
DEHP	117-81-7	17.79	149, 167, 57, 71	149	390.56
DINP	28553-12-0	20.0-21.3	149, 71, 57, 293	149	418.61

Table 4. Typical standard curves used for phthalate determination. Evaluation of linearity (R^2), limit of detection (LOD) and limit of quantification (LOQ).

Chemical	Minimum level (mg/L)	Maximum level (mg/L)	Slope	Intercept origin	R^2	LOD (mg/L)	LOQ (mg/L)
DEP	0.053	1.70	182525	-2775.8	0.99771	0.005	0.013
DBP	0.058	1.84	365291	-40707	0.99448	0.004	0.01
BBP	0.056	1.80	96154	-4398.4	0.99512	0.01	0.05
DEHP	0.055	1.76	183410	542.73	0.99390	0.01	0.03
DINP	0.631	20.2	8959.6	496.43	0.99775	0.4	0.99

Table 5. Number of phthalates detected in the tequila samples.

Year	0	1	2	3	4	5	Total
2013	0	2	13	0	0	0	15
2014	31	37	39	11	6	0	124
2015	20	37	5	2	0	0	64
2016	4	14	18	0	0	0	36
2017	10	20	12	0	0	0	42
2018	0	2	12	0	0	0	14
Total	65	112	99	13	6	0	295

Table 6. Summary of tequila samples contaminated with phthalates.

Chemical	Samples contained phthalate	Min (mg/kg)	Max (mg/kg)	Samples out of norm	% of samples out of norm
DEP	11	0.13	0.27	0	0
DBP	96	0.01	2.20	10	3.39
BBP	37	0.05	2.91	0	0
DEHP	224	0.03	4.64	15	5.08
DINP	5	1.64	3.43	0	0








Fig.1




Fig. 2.




Fig. 3.